arXiv:1610.05742v1 [math.CA] 15 Oct 2016

Product measures without utilizing integrals

Hasan Gül* and Süleyman Önal†

13.05.2016

Abstract: We show that product measures can be handled without resorting to integrals.

MSC: 28A35
Keywords: Analysis; Classical measure theory; Measures and integrals in product spaces

1 Introduction

In many textbooks on Real Analysis or Measure Theory, if not in all, product measures on product spaces are handled by utilizing the monotone and dominated convergence theorems which involve integrals with respect to some measures (cf. [1, 2, 3, 4, 5]). Of course, we do not find this approach to be circular, but feel a certain uneasiness.

The necessary foundation for studying a measure space (X, Σ, μ) is the characterization of null sets with respect to μ and the establishment of σ-additivity of it on the elements of the semiring (σ-algebra) Σ .

In this short note, we show that an introduction to product measure and laying the aforementioned foundation without resorting to integrals is possible in a very simple way.

2 Preliminaries

In this section, we recall some basic definitions.

Let X be a nonempty set.

Definition 2.1. A collection Σ of subsets of X is called a **semiring** if it satisfies the following properties:

1. $\emptyset \in \Sigma$,
2. If $A, B \in \Sigma$, then $A \cap B \in \Sigma$,

*Middle East Technical University, e032209@metu.edu.tr
†Middle East Technical University, osul@metu.edu.tr

3. For every $A, B \in \Sigma$, there exist $C_1, ..., C_n \in \Sigma$ such that $A \setminus B = \cup_{i=1}^{n} C_i$ and $C_i \cap C_j = \emptyset$ if $i \neq j$.

Definition 2.2. A nonempty collection Σ of subsets of X which is closed under finite intersections and complementation is called an **algebra** of sets. An algebra Σ of subsets of X is called a **σ-algebra** if every union of a countable collection of elements of Σ is again in Σ.

Definition 2.3. Let Σ be a semiring of subsets of X. A set function $\mu : \Sigma \to [0, \infty]$ is called a **measure** on Σ if it satisfies the following properties:

1. $\mu(\emptyset) = 0$,
2. If $\{A_n : n \in \mathbb{N}\}$ is a disjoint sequence in Σ satisfying $\cup_{n=1}^{\infty} A_n \in \Sigma$, then

$$\mu(\cup_{n=1}^{\infty} A_n) = \Sigma_{n=1}^{\infty} \mu(A_n)$$

holds.

Definition 2.4. A triplet (X, Σ, μ), where Σ is a semiring of subsets of X and μ is a measure on Σ is called a **measure space** .

Definition 2.5. Let (X, Σ, μ) be a measure space and $A \subseteq X$. If $\mu(A) < \infty$, then A is said to be of **finite measure**. If $A = \bigcup_{n \in \mathbb{N}} A_n$, where each A_n is of finite measure, then A is said to be of **σ-finite measure**.

Definition 2.6. A set function $\mu : P(X) \to [0, \infty]$ defined on the power set $\mathcal{P}(X)$ of X is called an **outer measure** if it satisfies these properties:

l. $\mu(\emptyset) = 0$,

2. If $A \subseteq B$, then $\mu(A) \leq \mu(B)$,

3. $\mu(\cup_{n=1}^{\infty} A_n) \leq \Sigma_{n=1}^{\infty} \mu(A_n)$ holds for every sequence $(A_n)_{n \in \mathbb{N}}$ of subsets of X.

Let $\mathcal{F}$ be a collection of subsets of X containing the empty set. Also, let $\mu : \mathcal{F} \to [0, \infty]$ be a set function such that $\mu(\emptyset) = 0$. For every subset A of X we define

$$\mu^*(A) = \inf\{\Sigma_{n=1}^{\infty} \mu(A_n) : (A_n) \text{ is a sequence of } \mathcal{F} \text{ with} A \subseteq \cup_{n=1}^{\infty} A_n\}.$$

If there is no sequence $(A_n)_{n \in \mathbb{N}}$ of $\mathcal{F}$ such that $A \subseteq \cup_{n=1}^{\infty} A_n$, then we let $\mu^*(A) = \infty$.

The set function $\mu^* : P(X) \to [0, \infty]$ is an outer measure (called the **outer measure generated** by the set function $\mu : \mathcal{F} \to [0, \infty]$) satisfying $\mu^*(A) \leq \mu(A)$ for each $A \in \mathcal{F}$.

Definition 2.7. Let (X, Σ_X, μ_X) and (Y, Σ_Y, μ_Y) be measurable spaces. The subset

$$\Sigma_X \otimes \Sigma_Y := \{A \times B : A \in \Sigma_X, B \in \Sigma_Y\}$$

of $X \times Y$ is called the **product semiring**. The set function $\mu_X \times \mu_Y : \Sigma_X \otimes \Sigma_Y \to [0, \infty]$ defined by

$$\mu_X \times \mu_Y(A \times B) = \mu_X(A) \cdot \mu_Y(B) \text{ for all } A \times B \in \Sigma_X \otimes \Sigma_Y.$$

is a measure on $\Sigma_X \otimes \Sigma_Y$ and is called the **product measure** of μ_X and μ_Y and is finitely additive.

3 Product measures

Theorem 1. *Let (X, Σ_X, μ_X) and (Y, Σ_Y, μ_Y) be measurable spaces where Σ_X, Σ_Y are semi-rings ($\sigma-$algebras) and μ_X, μ_Y are $\sigma-$ additive measures. Let $D \subseteq X \times Y$ and suppose $D \subseteq \cup_{n\in\mathbb{N}} B_n \times C_n$ where $\{B_n \times C_n : n \in \mathbb{N}\}$ are pairwise disjoint sequences with $B_n \in \Sigma_X$ and $C_n \in \Sigma_Y$. Define $D^{>r} := \{x \in X : \mu_Y^*(D^x) > r\}$ for $r \in \mathbb{R}^+$ and $D^x := \{y : (x, y) \in D\}$. Let μ_X^*, μ_Y^* be outer measures generated by μ_X and μ_Y, respectively, and suppose $\mu_X^*(D^{>r}) > s$ for some $s \in \mathbb{R}^+$, Then, there exists a finite subset $F \subseteq \mathbb{N}$ such that*

$$rs < \sum_{n\in F} \mu_X(B_n)\mu_Y(C_n)$$

.

Proof. Let $x \in D^{>r}$ and define $N_x := \{n \in \mathbb{N} : x \in B_n\}$. Then, $D^x \subseteq \bigcup_{n\in N_x} C_n$. Now

$$r < \mu_Y^*(D^x) \leq \Sigma_{n\in N_x}\mu_Y^*(C_n) \leq \Sigma_{n\in N_x}\mu_Y(C_n).$$

Let M_x be a finite subset of $N_x \subseteq \mathbb{N}$ such that $r < \Sigma_{n\in M_x}\mu_Y(C_n)$. Note that $x \in B_n$ for all $n \in M_x$. Thus, $D^{>r} \subseteq \bigcup_{x\in D^{>r}}(\bigcap_{n\in M_x} B_n)$ and the family $\{\bigcap_{n\in M_x} B_n : x \in D^{>r}\} \subseteq \Sigma_X$ is countable since $\{M_x : M_x \subseteq \mathbb{N}\} \subseteq \mathcal{P}_{<\aleph_o}(\mathbb{N})$.

We have

$$s < \mu_X^*(D^{>r}) \leq \mu_X^*(\bigcup_{x\in D^{>r}} (\bigcap_{n\in M_x} B_n)).$$

Thus, there exist a finite subset M of $D^{>r}$ with

$$s < \mu_X^*(\bigcup_{x\in M} (\bigcap_{n\in M_x} B_n).$$

Set $F := \bigcup_{x\in M} M_x$. Since $\bigcup_{x\in M} (\bigcap_{n\in M_x} B_n \times \bigcup_{n\in M_x} C_n) \subseteq \bigcup_{n\in F} (B_n \times C_n)$ we have $rs < \sum_{n\in F} \mu_X(B_n)\mu_Y(C_n)$ by the finite additivity of $\mu_X \times \mu_Y$ on the semiring $\Sigma_X \otimes \Sigma_Y$.

□

Corollary 1. *Let (X, Σ_X, μ_X) and (Y, Σ_Y, μ_Y) be measurable spaces where Σ_X, Σ_Y are semi-rings ($\sigma-$algebras) and μ_X and μ_Y are $\sigma-$ additive measures. Then $\mu_X \times \mu_Y$ is countably additive on $X \times Y$.*

Proof. Let (X, Σ_X, μ_X) and (Y, Σ_Y, μ_Y) be measurable spaces where Σ_X , Σ_Y are semi-rings ($\sigma-$algebras) and μ_X, μ_Y are $\sigma-$ additive measures. Let $D = B \times C = \bigcup_{n\in\mathbb{N}}(B_n \times C_n)$ with $B_n, B \in \Sigma_X$ and $C_n, C \in \Sigma_Y$ for all $n \in \mathbb{N}$ and $B_n \times C_n$ are pairwise disjoint for all $n \in \mathbb{N}$.

We may suppose, without loss of generality, $\mu(B) > 0$ and also,$\mu(C) > 0$.

Suppose $t < \mu_X(B) \cdot \mu_Y(C)$. We may find $r, s \in \mathbb{R}^+$ with $0 < r < \mu_X(B)$ and $0 < s < \mu_Y(C)$ such that $t < r \cdot s$. Consider $(B \times C)^x$ where $x \in B$. Clearly, $\mu^*((B \times C)^x) = \mu(C)$ and $D^{>r} = B$. Thus, $r < \mu(C)$ and $s < \mu^*(D^{>r})$. By the theorem above, we may find a finite subset F of $\mathbb{N}$ such that

$$t < \sum_{n \in F} \mu(A_n) \cdot \mu(B_n).$$

□

Corollary 2. *Let (X, Σ_X, μ_X) and (Y, Σ_Y, μ_Y) be measurable spaces where Σ_X, Σ_Y are semi-rings ($\sigma-$algebras) and μ_X, μ_Y are $\sigma-$ additive measures. If $D \subseteq X \times Y$ and $(\mu_X \times \mu_Y)^*(D) = 0$ then $\mu_Y^*(D^x) = 0$ for μ_X-almost all $x \in X$. Conversely, if D is $(\mu_X \times \mu_Y)^*$ measurable and $\mu_Y^*(D^x) = 0$ for μ_X-almost all x, then $(\mu_X \times \mu_Y)^*(D) = 0$, provided μ_X, μ_Y are σ-finite measures.*

Proof. If $D \subseteq X \times Y$ and $(\mu_X \times \mu_Y)^*(D) = 0$ then $\mu_X^*(D^{>r}) = 0$ for all $r > 0$ by the theorem. Hence $\mu_X^*(\bigcup_{n \in \mathbb{N}} D^{>\frac{1}{n}}) = 0$. Thus, $\mu_Y^*(D^x) = 0$ for μ_X-almost all $x \in X$.

Conversely, let μ_X, μ_Y be σ-finite measures. Suppose D is $(\mu_X \times \mu_Y)^*$ measurable and for μ_X-almost all x, $\mu_Y^*(D^x) = 0$. Then $(\mu_X \times \mu_Y)^*(D) = 0$.

By the σ-additivity of measures, we may assume $\mu_X(X)$, $\mu_Y(Y) < \infty$. Suppose $\mu_Y(D^x) = 0$ for μ_X-almost all $x \in X$. Since $D^c = X \times Y \setminus D$, we have $\mu_Y((D^c)^x) = \mu_Y(Y)$ for μ_X-almost all $x \in X$. Thus,

$$\mu_X^*(\{x : \mu_Y((D^c)^x) = \mu_Y(Y)\}) = \mu_X(X)$$

Hence, by the theorem, $(\mu_X \times \mu_Y)^*(D^c) = \mu_X(X) \cdot \mu_Y(Y)$. Therefore, $(\mu_X \times \mu_Y)^*(D) = 0$.

Note that both monotone and dominated convergence theorems themselves may be proved with this approach, which is, using measure theory instead of integration with a slightly different interpretation.

References

[1] C. D. ALIPRANTIS and O. BURKINSHAW, Principles of Real Analysis, 3rd ed., Academic Press, New York, 1998.

[2] E. HEWITT and K. STROMBERG, Real and Abstract Analysis, Springer-Verlag, New York, 1965.

[3] S. LANG, Real and Functional Analysis, Springer-Verlag, New York, 1993.

[4] H. ROYDEN, Real Analysis, Macmillan, New York, 1988.

[5] W. RUDIN, Real and Complex Analysis, McGraw-Hill, New York, 1966.

□